

## **TB cough aerosol analysis in a Peruvian shantytown**

P. Selvam\*, B. Herrera\*, RH. Gilman, J. Alvarado, E. Ramos, T. Valencia, S. Carrera, J. Alva, J. Franco, M. Haro, R. Montoya, CA. Evans

Universidad Peruana Cayetano Heredia, Lima, Peru; Asociacion Benefica PRISMA, Lima, Peru; Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

**BACKGROUND:** This study evaluated a novel diagnostic technique that utilizes a selective solid culture medium and a direct patient cough aerosol sample. Using a cough sample obviates sputum decontamination, may indicate infectiousness and has potential utility for patients who have difficulty expectorating sputum.

**METHODS:** Suspected tuberculosis patients (n=399) were asked to cough ten times on a cough plate for direct culture. For comparison, conventional sputum samples from each participant were decontaminated and tested for tuberculosis by microscopy and culture in Middlebrook 7H9 broth.

**RESULTS:** 45.6% (180/395) of subjects were sputum culture positive. Compared with conventional sputum culture, the 164 uncontaminated cough plates had sensitivity of 14% (23/164, 95%CI: 9.1–20.3). Cough plate sensitivity tended to be higher for subjects with strongly positive sputum microscopy indicating high bacterial load [sensitivity 31.0% (9/29, 95%CI: 15.3 – 50.8)] but five subjects had positive sputum and cough plate cultures despite negative sputum microscopy.

**CONCLUSION:** This novel cough plate test for tuberculosis identified a minority of sputum culture-positive patients. This cohort of 180 tuberculosis patients is being followed prospectively to determine whether this simple test for tuberculosis aerosolization predicts infectiousness for household contacts.

Presented by *Prabu Selvam* at the Johns Hopkins University Delta Omega public health research completion in March 2009 in Baltimore, USA and awarded third prize.